(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 63097, Nov. 2, 1979; 45 FR 1019, Jan. 4, 1980, as amended at 52 FR 18691, May 19, 1987; 52 FR 25212, July 6, 1987; 52 FR 36023, Sept. 25, 1987; 53 FR 32610, Aug. 26, 1988; 53 FR 40728, Oct. 18, 1988; 54 FR 41442, Oct. 10, 1989; 55 FR 8462, Mar. 8, 1990; 56 FR 14642, Apr. 11, 1991, 56 FR 16002, Apr. 19, 1991; 59 FR 31139, June 17, 1994; 62 FR 35076, June 30, 1997; 62 FR 38906, July 21, 1997; 65 FR 45877, July 26, 2000; 66 FR 22117, May 3, 2001; 68 FR 36913, June 20, 2003]

§ 522.1696 Penicillin G procaine implantation and injectable dosage forms.

§522.1696a Penicillin G benzathine and penicillin G procaine sterile suspension.

- (a) *Specifications.* Each milliliter of aqueous suspension contains penicillin G benzathine and penicillin G procaine, each equivalent to 150,000 units of penicillin G.
- (b) Sponsors. See sponsors in $\S510.600$ (c) of this chapter for the conditions of use in paragraph (d) of this section as follows:
- (1) Nos. 000008, 000856, 010515, 049185, 061623 for use as in paragraph (d)(1) of this section.
- (2) Nos. 000856 and 049185 for use as in paragraphs (d)(2)(i), (d)(2)(ii)(A), and (d)(2)(iii) of this section.
- (3) Nos. 010515, 059130, and 061623 for use as in paragraphs (d)(2)(i), (d)(2)(ii)(B), and (d)(2)(iii) of this section.
- (c) Related tolerances. See §556.510 of this chapter.
- (d) Conditions of use—(1) Horses, dogs, and beef cattle—(i) Amount—(A) Beef cattle. 2 milliliters per 150 pounds of body weight intramuscularly or subcutaneously. Repeat dosage in 48 hours.
- (B) *Horses.* 2 milliliters per 150 pounds of body weight intramuscularly. Repeat dosage in 48 hours.
- (C) *Dogs.* 1 milliliter per 10 to 25 pounds of body weight intramuscularly or subcutaneously. Repeat dosage in 48 hours.
- (ii) *Conditions of use.* Treatment of bacterial infections susceptible to penicillin G.
- (iii) *Limitations.* In beef cattle, treatment should be limited to two doses.

Not for use in beef cattle within 30 days of slaughter. Do not use in horses intended for food purposes. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) Beef cattle—(i) Amount. 2 milliliters per 150 pounds of body weight subcutaneously. Repeat dosage in 48

nours.

- (ii) Conditions of use. (A) Treatment of bacterial pneumonia (Streptococcus spp., Corynebacterium pyogenes, Staphylococcus aureus); upper respiratory infections such as rhinitis or pharyngitis (Cpyogenes); blackleg (Clostridium chauvoei).
- (B) As in paragraph (d)(2)(ii)(A) of this section; and prophylaxis of bovine shipping fever in 300- to 500-pound beef calves
- (iii) *Limitations*. Limit treatment to two doses. Not for use within 30 days of slaughter.

[66 FR 711, Jan. 4, 2001, as amended at 68 FR 34534. June 10. 2003]

§ 522.1696b Penicillin G procaine aqueous suspension.

- (a) *Specifications*. Each milliliter contains penicillin G procaine equivalent to 300,000 units of penicillin G.
- (b) *Sponsors*. See sponsor numbers in §510.600(c) of this chapter as follows:
- (1) Nos. 010515, 053501, and 059130 for use as in paragraph (d) of this section.
- (2) Nos. 055529 and 061623 for use as in paragraph (d)(2) of this section.
- (c) Related tolerances. See §556.510 of this chapter.
- (d) Conditions of use—(1) Dogs and cats—(i) Amount. 10,000 units per pound body weight daily by intramuscular injection at 24-hour intervals. Continue treatment at least 48 hours after symptoms disappear.
- (ii) *Indications for use*. Treatment of infections caused by penicillin-sensitive organisms.
- (iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Cattle, sheep, swine, and horses—(i) Amount. 3,000 units per pound body weight (1 milliliter per 100 pounds body weight) daily by intramuscular injection.
- (A) For Nos. 010515, 053501, 059130, and 061623: Continue treatment at least 48 hours after symptoms disappear.